

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12Q 1/70, A01N 43/04, A61K 31/70</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/44492</b> <b>(43) International Publication Date:</b> 27 November 1997 (27.11.97)
<b>(21) International Application Number:</b> PCT/US97/08686 <b>(22) International Filing Date:</b> 22 May 1997 (22.05.97) <b>(30) Priority Data:</b> 08/652,074 23 May 1996 (23.05.96) US <b>(71) Applicant:</b> THE ROCKEFELLER UNIVERSITY [US/US]; 1230 York Avenue, New York, NY 10021 (US). <b>(72) Inventors:</b> HALL, William, W.; 1230 York Avenue, New York, NY 10021 (US). TAKAHASHI, Hidehiro; Toyama, 1-23-1, Chinjuku-ku, Tokyo 162 (JP). <b>(74) Agents:</b> CORUZZI, Laura, A. et al.; Pennie & Edmonds, LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).		<b>(81) Designated States:</b> AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> TREATMENT OF HIV INFECTION BY INTERFERING WITH HOST TOPOISOMERASE INTERACTION WITH HIV GAG AND RT  <b>(57) Abstract</b>  The present invention relates to therapeutic protocols and pharmaceutical compositions designed to target topo (I) for the treatment of HIV infection. The invention relates to therapeutic modalities and pharmaceutical compositions for the treatment of HIV-infection using human topo (I) and its interaction with HIV gag and RT as a target for intervention. The invention further relates to the use of human topo (I) to enhance the activity of RT. The present invention also relates to the expression of human topo (I) in transgenic animals, in particular mice, as a system to study the HIV life cycle and the screen agents for their ability to interfere with the HIV life cycle.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TREATMENT OF HIV INFECTION BY INTERFERING WITH  
HOST TOPOISOMERASE INTERACTION WITH HIV GAG AND RT

This application is a continuation-in-part of co-pending application Serial No. 08/652,074, filed May 23, 1996, which  
5 is incorporated by reference herein in its entirety.

1. INTRODUCTION

The present invention relates to therapeutic protocols and pharmaceutical compositions designed to target  
10 topoisomerase I (topo I) as a treatment for HIV-infection. In a preferred embodiment, the present invention relates to therapeutic protocols designed to interfere with the interaction of topo I with the HIV proteins gag and reverse transcriptase (RT), as a treatment for HIV-infection. The  
15 present invention still further relates to transgenic animals which express human topo I, in particular transgenic mice, and their use as a system to screen agents for their ability to interfere with the HIV life cycle.

20

2. BACKGROUND OF THE INVENTION

2.1. THE HUMAN IMMUNODEFICIENCY VIRUS

The human immunodeficiency virus (HIV) has been implicated as the primary cause of the slowly degenerative immune system disease termed acquired immune deficiency  
25 syndrome (AIDS) (Barre-Sinoussi, F. et al., 1983, Science 220:868-870; Gallo, R. et al., 1984, Science 224:500-503). There are at least two distinct types of HIV: HIV-1 (Barre-Sinoussi, F. et al., 1983, Science 220:868-870; Gallo R. et al., 1984, Science 224:500-503) and HIV-2 (Clavel, F. et al.,  
30 1986, Science 233:343-346; Guyader, M. et al., 1987, Nature 326:662-669). Further, a large amount of genetic heterogeneity exists within populations of each of these types. In humans, HIV replication occurs prominently in CD4<sup>+</sup> T lymphocyte populations, and HIV infection leads to  
35 depletion of this cell type and eventually to immune incompetence, opportunistic infections, neurological dysfunctions, neoplastic growth, and ultimately death..

HIV is a member of the lentivirus family of retroviruses (Teich, N. et al., 1984, RNA Tumor Viruses, Weiss, R. et al., eds., CSH-Press, pp. 949-956). Retroviruses are small enveloped viruses that contain a single-stranded RNA genome, and replicate via a DNA intermediate produced by a virally-encoded reverse transcriptase, an RNA-dependent DNA polymerase (Varmus, H., 1988, Science 240:1427-1439). Other retroviruses include, for example, oncogenic viruses such as human T cell leukemia viruses (HTLV-I, -II, -III), and feline leukemia virus.

The HIV viral particle comprises a viral core, composed in part of capsid proteins, together with the viral RNA genome and those enzymes required for early replicative events. Myristylated Gag protein forms an outer shell around the viral core, which is, in turn, surrounded by a lipid membrane envelope derived from the infected cell membrane. The HIV envelope surface glycoproteins are synthesized as a single 160 Kd precursor protein which is cleaved by a cellular protease during viral budding into two glycoproteins, gp41 and gp120. gp41 is a transmembrane glycoprotein and gp120 is an extracellular glycoprotein which remains non-covalently associated with gp41, possibly in a trimeric or multimeric form (Hammarskjold, M. & Rekosh, D., 1989, Biochem. Biophys. Acta 989:269-280).

HIV is targeted to CD4<sup>+</sup> cells because a CD4 cell surface protein (CD4) acts as the cellular receptor for the HIV-1 virus (Dalglish, A. et al., 1984, Nature 312:763-767; Klatzmann et al., 1984, Nature 312:767-768; Maddon et al., 1986, Cell 47:333-348). Viral entry into cells is dependent upon gp120 binding the cellular CD4 receptor molecules (McDougal, J.S. et al., 1986, Science 231:382-385; Maddon, P.J. et al., 1986, Cell 47:333-348), explaining HIV's tropism for CD4<sup>+</sup> cells, while gp41 anchors the envelope glycoprotein complex in the viral membrane. While these virus:cell interactions are necessary for infection, there is evidence that additional virus:cell interactions are also required.

Purified HIV virions have been reported to have topoisomerase activity (Priel et al., 1990, EMBO J. 9:4167-4172). Topoisomerase I is an enzyme that modifies the topological state of the DNA by breakage and rejoining of  
5 single DNA strands and relaxing both positive and negative supercoils. In addition to playing a role in normal cell function, topo I of cellular origin has also been shown to be involved in the replication of a number of DNA viruses, where it has been associated with DNA replication, transcription,  
10 and integration (Wang et al. 1990, J. Virol. 64:691-699; Shaack et al., 1990, Nucleic Acids Research 18:1499-1508; Kawanishi 1993, J. Gen. Virology 74:2263-2268). It has been suggested that the topo I activity associated with HIV virions is viral in origin in that it differs from cellular  
15 topo I in its requirements for optimal enzyme activity (Priel et al., 1990, EMBO J. 9:4167-4172).

## 2.2. HIV TREATMENT

HIV infection is pandemic and HIV-associated diseases  
20 represent a major world health problem. Although considerable effort is being put into the design of effective therapeutics, currently no curative anti-retroviral drugs against AIDS exist. In attempts to develop such drugs, several stages of the HIV life cycle have been considered as  
25 targets for therapeutic intervention (Mitsuya, H. et al., 1991, FASEB J. 5:2369-2381). Many viral targets for intervention with HIV life cycle have been suggested, as the prevailing view is that interference with a host cell protein would have deleterious side effects. For example, virally  
30 encoded reverse transcriptase has been one focus of drug development. A number of reverse-transcriptase-targeted drugs, including 2',3'-dideoxynucleoside analogs such as AZT, ddI, ddC, and d4T have been developed which have been shown to be active against HIV (Mitsuya, H. et al., 1991, Science  
35 249:1533-1544). While beneficial, these nucleoside analogs are not curative, probably due to the rapid appearance of drug resistant HIV mutants (Lander, B. et al., 1989, Science

243:1731-1734). In addition, the drugs often cause toxic side effects such as bone marrow suppression, vomiting, and liver function abnormalities.

Attempts are also being made to develop drugs which can inhibit viral entry into the cell, the earliest stage of HIV infection. Here, the focus has thus far been on CD4, the cell surface receptor for HIV. Recombinant soluble CD4, for example, has been shown to inhibit infection of CD4<sup>+</sup> T cells by some HIV-1 strains (Smith, D.H. et al., 1987, Science 238:1704-1707). Certain primary HIV-1 isolates, however, are relatively less sensitive to inhibition by recombinant CD4 (Daar, E. et al., 1990, Proc. Natl. Acad. Sci. USA 87:6574-6579). In addition, recombinant soluble CD4 clinical trials have produced inconclusive results (Schooley, R. et al., 1990, Ann. Int. Med. 112:247-253; Kahn, J.O. et al., 1990, Ann. Int. Med. 112:254-261; Yarchoan, R. et al., 1989, Proc. Vth Int. Conf. on AIDS, p. 564, MCP 137).

The late stages of HIV replication, which involve crucial virus-specific processing of certain viral encoded proteins, have also been suggested as possible anti-HIV drug targets. Late stage processing is dependent on the activity of a viral protease, and drugs are being developed which inhibit this protease (Erickson, J., 1990, Science 249:527-533). The clinical outcome of these candidate drugs is still in question.

Attention is also being given to the development of vaccines for the treatment of HIV infection. The HIV-1 envelope proteins (gp160, gp120, gp41) have been shown to be the major antigens for anti-HIV antibodies present in AIDS patients (Barin, et al., 1985, Science 228:1094-1096). Thus far, therefore, these proteins seem to be the most promising candidates to act as antigens for anti-HIV vaccine development. Several groups have begun to use various portions of gp160, gp120, and/or gp41 as immunogenic targets for the host immune system. See for example, Ivanoff, L. et al., U.S. Pat. No. 5,141,867; Saith, G. et al., WO 92/22,654; Shafferman, A., WO 91/09,872; Formoso, C. et al., WO



90/07,119. To this end, vaccines directed against HIV proteins are problematic in that the virus mutates rapidly rendering many of these vaccines ineffective. Clinical results concerning these candidate vaccines, however, still remain far in the future.

Thus, although a great deal of effort is being directed to the design and testing of anti-retroviral drugs, effective, non-toxic treatments are still needed.

10

### 3. SUMMARY OF THE INVENTION

The present invention relates to therapeutic protocols and pharmaceutical compositions designed to target topo I for the treatment of HIV infection. The invention relates to therapeutic modalities and pharmaceutical compositions for the treatment of HIV-infection using human topo I and its interaction with HIV gag and RT as a target for intervention.

The present invention relates to animal cell lines expressing human topo I, in particular mouse cell lines, and their use as a system to study the HIV life cycle and screen agents for their ability to interfere with the HIV life cycle. The present invention also relates to human topo I transgenic animals, in particular mice, and their use as a system to study the HIV life cycle and to screen agents for their ability to interfere with the HIV life cycle.

25 The invention is based, in part, on the Applicants' surprising discoveries that (1) human topo I interacts with and is activated by HIV gag in a species specific manner; (2) the interaction between human topo I and gag is required to enhance HIV RT activity; and (3) the interactions between human topo I and HIV gag and RT are required to support HIV replication. This model is based on the Applicants' observation that murine cells expressing human CD4 are not able to support HIV replication. However, murine cells expressing both human CD4 and human topo I are able to support HIV replication. The expression of human topo I was also shown to enhance the activity of HIV RT in murine cells.

That human topo I interacts with and is activated by HIV gag, and that this complex is required for activation of HIV RT is further supported by the working examples described *infra* which demonstrate (1) that gag proteins activate  
5 cellular topo I and immunoprecipitated-gag proteins induce topo I activity in a species specific manner; (2) mouse cells expressing both human CD4 and topo I and infected with HIV, effectively reverse transcribe the HIV RNA genome; and (3) the topo I inhibitor, TAN134A, which attacks the topo I site  
10 directly, inhibits HIV RT activity in murine cells.

The invention further relates to a murine model for HIV replication, in which transgenic mice expressing both the human CD4 cell surface protein and human topo I are able to support HIV-1 replication. The present invention also  
15 encompasses a murine model for HIV replication, in which transgenic mice express human topo I and a HIV pseudovirus is used to infect the animals. The HIV pseudovirus may contain an envelope protein from a virus with a natural tropism for murine cells, such as the murine leukemia virus, which  
20 bypasses internalization of the HIV virus by the murine cells. These transgenic mice have utility to screen for other host cellular components required to support the HIV life cycle (i.e., entry, replication and assembly), in addition to screen for drugs and compounds which may have  
25 anti-HIV activity.

The invention relates to various modalities of treatment for HIV infection based on the proposed model. The invention further relates to the use of the murine HIV model system for screening test compounds, such as drugs, ligands (natural or  
30 synthetic), proteins, peptides and small organic molecules for their ability to interfere with the interaction between human topo I and HIV gag and RT.

The present invention further relates to the use of such identified inhibitors in pharmaceutical compositions designed  
35 to inhibit human topo I and/or the interaction between human topo I and HIV gag and/or HIV RT for the treatment and/or prevention of HIV infection. The present invention further

encompasses the preparation of such pharmaceutical compositions for the treatment and/or prevention of HIV infection.

The invention also encompasses combinations of a  
5 topoisomerase I inhibitor with a least one other antiviral having a different site of action than the viral replication inhibitor. Such a combination provides an improved therapy based on the dual action of these therapeutics whether the combination is synergistic or additive.

10

### 3.1. DEFINITIONS

As used herein, the term "topoisomerase I" or "topo I" refers to the protein topoisomerase I which has the activity of modifying the topological state of DNA, and any derivative  
15 of topo I thereof, or fragments or peptides having an amino acid sequence corresponding to topo I.

As used herein, the term "HIV pseudovirus" refers to a HIV virus that expresses a coat protein or an envelope protein which renders the virus capable of infecting a mouse  
20 cell. An example of such a protein is the envelope protein of the murine leukemia virus.

As used herein, the term "to target" means to inhibit, block, or prevent gene expression, enzymatic activity, or interaction with other cellular or viral factors.

25 As used herein, the term "treating or preventing HIV infection" means to inhibit the replication of the HIV virus, to inhibit HIV transmission, or to prevent HIV from establishing itself in its host, and to ameliorate or alleviate the symptoms of the disease caused by HIV  
30 infection. The treatment is considered therapeutic if there is a reduction in viral load, decrease in mortality and/or morbidity.

As used herein, the term "pharmaceutically acceptable carrier" refers to a carrier medium that does not interfere  
35 with the effectiveness of the biological activity of the active ingredient, is chemically inert and is not toxic to the patient to whom it is administered.

The term "therapeutic agent" refers to any molecule compound or treatment, preferably an antiviral, that assists in the treatment of a viral infection or the diseases caused thereby.

5

#### 4. BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1. Southern hybridization analysis of transfected supercoiled plasmid DNA in HeLa cells infected with RVV expressing HIV Pr55gag (RVV-Pr55gag) or wild-type vaccinia virus (wild-type VV). OC refers to open circle and relaxed DNA, and SC refers to supercoiled DNA.

FIGURE 2. Topo I activity after the interactions of human topo I with GST-Pr55gag or GST alone. Five-tenths microgram of supercoiled plasmid DNA was incubated with 0.1 unit of human topo I and GST-Pr55gag or GST alone at final concentrations of 50, 10, 2, 0.4, 0.08  $\mu\text{g/ml}$  (+) for 20 minutes at 37°C. Reactions without human topo I are indicated (-). Products were analyzed by electrophoresis on 1% agarose gels.

FIGURE 3. Influence of cellular topo I on HIV-1 RT activity. FIGURE 3A. Schematic representation of the RNA template employed in the RT assays. The 5'/3' RNA contains the 5'R, U5, and part of the gag region truncated to the nef gene and the 3' U3, and poly(A) tail (PA). An end-labeled primer (\*) was employed to initiate cDNA synthesis and would be expected to result in a product of 1481 nucleotides (nt). FIGURE 3B. Phosphorescence image analysis of cDNA products following agarose gel electrophoresis. The presence or absence of topo I is indicated by + and -, respectively. Similarly the presence or absence of CPT is indicated by + and -.

FIGURE 4. Analysis of ssDNA synthesis and strand transfer. FIGURE 4A. Schematic representation of two RNA templates employed. The 5' RNA represents the 5' end of HIV-1 RNA and contains the R, U5, and PBS. The 3' RNA represents the 3' end of the HIV-1 RNA and contains the R region and the

poly(A) tail. ssDNA synthesis was initiated by an end labeled primer (\*) corresponding to the primer binding site (PBS) in the 5' RNA. After strand transfer to the 3'RNA, this would be expected to result in a cDNA product of 867 nucleotides. FIGURE 4B. Phosphorescence image analysis of the cDNA product after agarose gel electrophoresis. The presence and absence of 5'RNA and 3'RNA templates are indicated by + and -, respectively. The effects of CPT at concentrations of 400, 40 and 4  $\mu$ M are indicated by +. Topo I was used at the final concentration of 0.16 unit per 50  $\mu$ l in all experiments.

FIGURE 5. Southern hybridization of DNA from HIV-1 infected cells with drug treatment. Mouse cells, MT2, infected with the HIV-1 IIIB strain were treated with AZT, CPT or TAN at concentrations (from right to left) AZT 0.01, 0.1, 1.0, 10  $\mu$ g/ml, CPT 0.01, 0.1, 1.0  $\mu$ g/ml, TAN 0.01, 0.1, 1.0  $\mu$ g/ml respectively.

FIGURE 6. Topo I activity after the interactions of human topo I or calf topo I with GST-Pr55gag. Five tenths microgram of supercoiled pUC19 DNA was incubated with 0.1 unit of human topo I or calf topo I and GST-Pr55gag at final concentrations of 50, 10, 2, 0.4, 0.08  $\mu$ g/ml (+) for 15 minutes at 37°C. Reactions without topo I are indicated as (-).

FIGURE 7. Establishment of murine cell lines stably expressing human CD4 and human topoisomerase I. FIGURE 7A. Selected cell lines were stained with Leu3a and FITC conjugated goat anti-mouse sera and analyzed by flow cytometry as indicated. FIGURE 7B. Expression of c-myc epitope tagging human topo I. The murine fibroblast cell line stably expressing CD4 was again stably transfected with control expression vector or c-myc epitope-tagging human topo I expression vector in the presence of Hygromycine resistant vector were analyzed by immunoblotting with rabbit anti-myc serum.

FIGURE 8. Investigation of the role of human topo I on HIV-1 reverse transcription in L929 cells. L929 cells were

infected with HIV-1 (IIIB) by incubation for 24 hrs. and following trypsinization, cells were incubated further 7 days. Proviral DNA synthesis was analyzed by polymerase chain reaction (PCR) techniques employing gag(c) or U3-U5(d) specific primers. PCR using the primers for gag-amplification, SK38, 5'ATAATCCACCTATCCCAGTAGGAGAAAT3' (1544-1571) and SK39, 5'TTTGGTCCTTGTCTTATGTCCAGAATGC3' (1631-1658) or U3-U5-amplification, HL30, 5'GACAGCCTCCTAGCATTTCGTCAC3' (265-288) and HL10, 5'AGGGTCTGAGGGATCTCTAG3' (588-607). It can be seen that in the cells expressing only human CD4 (L929/hCD4) provirus could not be detected. In contrast, in the cells expressing both human CD4 and complete topo I (L929/hCD4/htopoI), provirus could be readily detected.

FIGURE 9. Recombinant plasmids for the production of HIV-1 pseudotype. FIGURE 9A. The recombinant HIV-1 vector. A 1.2 kb deletion in the env gene of pNL43 was made, leaving the rev responsive element and tat and rev exons intact. SV40 ori and hygromycin or brastcydine resistant sequences were inserted into the env deletion site. These constructs were named as HIV-hyg or HIV-BSD respectively. FIGURE 9B. Structure of the MLV ecotropic env expression vector. Transcription is driven by SR<sub>g</sub> promoter.

FIGURE 10. Immunoblot analysis of gag proteins expressed by COS1 cells transfected with HIV-hyg. When HIV-hyg or HIV-bsd and MLV-eco were cotransfected into COS-1 cells, bands corresponding to the processed (p24) and precursor form (p55) of HIV-1 gag proteins were observed.

FIGURE 11. Nucleotide sequence of human topoisomerase I Genebank Accession Number N83271.

FIGURE 12. Construction of plasmids expressing human CD4 or human topoisomerase I under the granzyme B promoter, which allows expression in peripheral blood cells and in particular NK cells and T lymphocytes. The p687 plasmid is derived from puc19. The hCD4 DNA was prepared by PCR amplification and inserted at the Bam HI site of p687. The Bgl II/ Bam HI fragment of the htopo I sequence (as described

in Fig. 11) was inserted at the Bam HI/ Bgl II site of p687. Both plasmids were linearized by digestion with SalI prior to microinjection.

FIGURE 13. Plasmid *plck-hGH* contains the *lck* promoter, 5 which allows expression in peripheral T lymphocytes. The plasmid contains one Bam HI cloning site. The Bam HI/ Bgl II fragment of the *htopo I* sequence (as described in Fig. 11) was inserted at the Bam HI site of *plck-hGH*. The *hCD4* DNA was prepared by PCR amplification and inserted at the Bam HI 10 site of *plck-hGH*.

#### 5. DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to therapeutic protocols and pharmaceutical compositions designed to target *topo I* for 15 the treatment of HIV infection. The invention relates to therapeutic modalities and pharmaceutical compositions for the treatment of HIV-infection using human *topo I* and its interaction with HIV *gag* and RT as a target for intervention. The invention further relates to the use of human *topo I* to 20 enhance the activity of RT.

The present invention relates to targeting human *topo I* and/or the interaction between human *topo I* and HIV proteins *gag* and RT as a means for treating HIV infection. The invention is based, in part, on the Applicants' discovery 25 of the specific interaction between human *topo I* and HIV proteins *gag* and RT and the requirement of this interaction to support HIV replication.

The present invention relates to therapeutic modalities and pharmaceutical compositions for the treatment of HIV 30 infection; these agents are designed to interfere with the interaction between human *topo I* and HIV proteins *gag* and RT. A variety of techniques and compositions may be utilized to inhibit the interaction of *topo I* and HIV proteins *gag* and RT, thereby inhibiting HIV-infection.

35 The invention further relates to a murine model for HIV replication, in which transgenic mice express human *topo I* alone or the CD-4 cell surface protein and human *topo I*.

These transgenic mice are able to support HIV life cycle (i.e., entry, replication and assembly). The transgenic mice of the present invention have utility as a screen to identify other host cellular components required to support the HIV life cycle. In a preferred embodiment of the invention these transgenic mice have utility in screening T cell genomic libraries to identify additional human proteins required to support the HIV life cycle. These transgenic mice also have utility as a screen for drugs and compounds which may have anti-HIV activity.

Another embodiment of the invention relates to small organic molecules that interfere with the human topo I or the interaction of human topo I and HIV proteins to inhibit HIV infection. The present invention additionally relates to screening assays to identify compounds which inhibit HIV infection, these compounds are selected for their activity in inhibiting human topo I or blocking the interaction between human topo I and HIV proteins gag and RT. Such identified compounds would have utility in the treatment and prevention of HIV-infection.

The therapeutic modalities of the invention further encompass combination therapy in which an agent which interferes with the interaction of human topo I and HIV protein and at least one other therapeutic agent are administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially, including cycling therapy. Cycling therapy involves the administration of a first antiviral compound for a period of time, followed by the administration of a second antiviral compound for a period of time and repeating this sequential administration, i.e., the cycle, in order to reduce the development of resistance to one of the therapies.

The novel antiviral combinations of the present invention provide a means of treatment which may not only reduce the effective dose of either drug required for antiviral activity, thereby reducing toxicity, but may also improve the absolute antiviral effect as a result of



attacking the virus through multiple mechanisms. Similarly, the novel antiviral combinations provide a means for circumventing the development of viral resistance to a single therapy, thereby providing the clinician with a more  
5 efficacious treatment.

Therapeutic agents to be used in combination with a topoisomerase I inhibitor encompass a wide variety of known treatments. Preferably, the combinations employ a topo I inhibitor in combination with agents with a different mode of  
10 attack. Such agents include but are not limited to: antivirals, such as cytokines, inhibitors of reverse transcriptase, inhibitors of viral capping, and inhibitors of viral protease.

The present invention still further relates to the  
15 Applicants' discovery that the ability of topo I to enhance the activity of endogenous RT can be utilized to improve the efficiency of gene therapy.

20 5.1. THE ROLE OF HUMAN TOPO I AND ITS  
INTERACTION WITH HIV GAG AND RT  
IN HIV INFECTION

The present invention is based, in part, on the Applicants' surprising discovery that (1) human topo I interacts with and is activated by HIV gag in a species specific manner; (2) the interaction between human topo I and  
25 gag is required to enhance HIV RT activity; and (3) the interactions between human topo I and HIV gag and RT are required to support HIV replication. This model is based on the Applicant's observation that although murine cells expressing human CD4 are not capable of supporting HIV  
30 replication (Maddon et al., 1986, Cell 47: 333-348), murine cells expressing both human CD4 and human topo I are able to support HIV replication. This discovery is exemplified in Sections 6, 7 and 8 *infra*, which demonstrate  
35 that the activation of human topo I by gag is species specific, in that the enhancement of calf or mouse topo I activity could not be detected in the presence of HIV gag,

even at high concentrations of gag. The expression of human topo I was shown to enhance the activity of HIV RT in murine cells.

That human topo I interacts with and is activated by HIV 5 gag, and that this complex is required for activation of HIV RT is further supported by the working examples described *infra* which demonstrate (1) that gag proteins activate cellular topo I in a species specific manner; (2) mouse cells expressing human CD4 and topo I infected with HIV effectively 10 reverse transcribe the HIV RNA genome; and (3) the topo I inhibitor, TAN134A, which attacks the topo I active site directly, inhibits RT activity in murine cells.

15 5.2. HUMAN TOPO I AS A TARGET FOR  
INTERVENTION IN THE TREATMENT OF  
HIV INFECTION


As discussed above, the Applicants' work shows that inhibition of human topo I or the interaction between human topo I and HIV proteins is effective in inhibiting cellular infection by HIV. A variety of techniques and compositions 20 may be utilized to target human topo I to inhibit its activity or to inhibit the interaction between human topo I and HIV proteins, thereby inhibiting HIV infection.

25 5.2.1. TOPOISOMERASE INHIBITORS

Compounds which may be used in accordance with the present invention encompass any compound which targets human topo I to inhibit its activity or interferes with the interaction between human topo I and its activation by HIV gag or its ability to activate HIV RT, including but not 30 limited to neutralizing antibodies against human topo I, or HIV proteins gag and RT. Other examples of compounds include but are not limited to peptides (such as, for example, peptides representing those regions of human topo I required for its interaction with HIV gag or RT), phosphopeptides, 35 small organic or inorganic molecules, or antibodies (including, for example, polyclonal, monoclonal, humanized,

anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')<sub>2</sub> and FAb expression library fragments, and epitope-binding fragments thereof). Techniques for determination of effective doses and administration of such compounds are 5 described below, in Section 5.6 *infra*.

Compounds which may be used in accordance with the present invention include, but are not limited to inhibitors of topoisomerase. Examples of topoisomerase inhibitors include, but are not limited to, camptothecin, synthetic and 10 semi-synthetic derivatives thereof, e.g., irinotecan, topotecan and derivatives thereof and pharmaceutically acceptable salts thereof. Camptothecin and derivatives thereof which may be used in accordance with the present invention are described in Jaxel et al. 1989 Cancer Research 15 49:1465-1469 and Giovanella et al. 1989 Science 246:1046-1048 both of which are incorporated herein by reference in their entireties. By the term "camptothecin" as used herein is meant the compound of the formula:



Substituents of the above shown structure which may be 35 used, in accordance with the invention, include as follows: hydroxy, nitro, amino, chloro, bromo, iodo, fluoro, C<sub>1-8</sub> alkyl, C<sub>1-8</sub> alkoxy, trifluoromethyl, aminomethyl, amido,


hydrazino, azido, formyl, and cyano groups as well as groups comprising amino acids bonded to aromatic ring via the amino-nitrogen atom. Preferred alkyl groups include methyl, ethyl, propyl, butyl, isopropyl, isobutyl, and sec-butyl groups.

5 Preferred alkoxy groups include methoxy, ethoxy, propoxy and isopropoxy groups.

Preferred camptothecin derivatives which may be used in accordance with the present invention include, but are not limited to, 11-methoxy-20(RS)-camptothecin, 11-methoxy-20(S)-  
10 camptothecin, 11-hydroxy-20(S)-camptothecin, 10-hydroxy-20(RS)-camptothecin, 10-hydroxy-20(S)-camptothecin, 9-methoxy-20(RS)-camptothecin, 9-methoxy-20(S)-camptothecin, 9-hydroxy-20(RS)-camptothecin, 9-hydroxy-20(S)-camptothecin, 10-nitro-20(RS)-camptothecin, 10-nitro-20(S)-camptothecin,  
15 10-amino-20(RS)-camptothecin, 10-amino-20(S)-camptothecin, 9-nitro-20(RS)-camptothecin, 9-nitro-20(S)-camptothecin, 9-amino-20(RS)-camptothecin, 9-amino-20(S)-camptothecin, 11-nitro-20(RS)-camptothecin, 11-nitro-20(S)-camptothecin, 11-amino-20(RS)-camptothecin, 11-amino-20(S)-camptothecin,  
20 10,11-dihydroxy-20(RS)-camptothecin, 10,11-dihydroxy-20(S)-camptothecin, 10-chloro-20(RS)-camptothecin, 10-chloro-20(S)-camptothecin, 10-methyl-20(S)-camptothecin, 11-formyl-20(RS)-camptothecin and 11-formyl-20(S)-camptothecin, and pharmaceutically acceptable salts thereof, including  
25 hydrochloride salt, acetate salt and methanesulfonic acid salt. Derivatives of camptothecin which may be used in accordance with the present invention and methods of synthesizing such derivatives are described in U.S. Patents 5,496,830, 5,364,858, 5,244,903, 5,004,758 and PCT  
30 publication WO95/09169 each of which is incorporated herein by reference in their entireties.

Water soluble derivatives of camptothecin may also be used in accordance with the invention. Topotecan is the most preferred compound of the water soluble camptothecin analog  
35 class.

By the term "topotecan" as used herein is meant the compound of the formula:




(S)-9-dimethylaminomethyl-10-hydroxycamptothecin and any  
15 pharmaceutically acceptable salt, hydrate or solvate thereof  
as described in WO 92/14470 incorporated herein by reference  
in its entirety. Topotecan's chemical name is (S)-  
10[(dimethylamino)methyl]-4-ethyl-4,9-dihydroxy-1H-  
pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-  
20 dione.

Topotecan is water-soluble by virtue of the presence of  
the basic side-chain at position 9 which forms salts with  
acids. Preferred salt forms of topotecan include the  
hydrochloride salt, acetate salt and methanesulfonic acid  
25 salt. An alkali metal salt form of the carboxylate formed on  
alkaline hydrolysis of the E-ring lactone of topotecan would  
also yield a soluble salt, such as the sodium salt.

The preparation of topotecan (including pharmaceutically  
acceptable salts, hydrates and solvates thereof) as well as  
30 the preparation of oral and parenteral pharmaceutical  
compositions comprising topotecan and an inert,  
pharmaceutically acceptable carrier or diluent, is  
extensively described in U.S. Patent Number 5,004,758. The  
same extensive description is found in European Patent  
35 Application Number 88311366.4, published on June 21, 1989 as  
Publication Number EP 0 321 122.

Examples of additional water soluble camptothecin analogs which may be used in accordance with the present invention include analogs of the following structure:



15 wherein: R1 and R2 are H, (C<sub>1-4</sub>) alkyl or may together form a 5 to 7 membered heterocyclic ring containing the nitrogen and up to one additional heteroatom such as O, S or NR<sup>3</sup> wherein R<sup>3</sup> is H or (C<sub>1-4</sub>) alkyl, and pharmaceutically acceptable salts thereof, including hydrochloride salt, acetate salt and

20 methanesulfonic acid salt (see PCT WO 9638449, published Dec 5, 1996, U.S. Patent 5,004,758, issued April 2, 1991, each of which is incorporated herein by reference in their entireties). The present invention also encompasses water soluble camptothecin derivatives which are substituted in the

25 7 position as described in U.S. Patent 5,559,235 as incorporated herein by reference in its entirety.

Semisynthetic derivatives of camptothecin, such as irinotecan and pharmaceutically acceptable salts thereof, including hydrochloride salt, acetate salt and

30 methanesulfonic acid salt, may also be used in accordance with the methods of the present invention. The preparation of irinotecan is described in U.S. Patent 4,604,463 incorporated herein by reference in its entirety.


Topoisomerase inhibitors also to be used in accordance

35 with the present invention, include, but are not limited to those of microbial origin such as TAN-1496 A, B, C, D and E, isolated from a fungal strain, TAN-1609, also known as

herboxidiene (see J. Takeda Research Labs, 1996, 55:149). Also to be used in accordance with the present invention are topoisomerase inhibitors TAN 134A, TAN 134B, TAN 1518A and TAN 1518B. (See, Horuguchi et al., 1995, Biochem Pharmac. 59:1395-1401 and Funabashi et al. 1994, J. Antibiotic 47:554-556 both of which are incorporated by reference herein in their entireties). Examples of the TAN compounds to be used in accordance with present invention may have the basic structure:

10

15



	n	R <sub>1</sub>	R <sub>2</sub>
A	2	Ac	Ac
B	2	Ac	H
C	3	Ac	Ac
D	2	H	H
E	4	Ac	Ac

20

In a preferred embodiment of the present invention, the protease inhibitor CAMPTOSAR, (Smith Kline Beecham, Philadelphia, PA) may be used in accordance with the present invention. CAMPTOSAR is a brand of irinotecan hydrochloride injection, a semisynthetic derivative of camptothecin. The chemical name is (4S)-4,11-diethyl-4-hydroxy-9-[(4-piperidinopiperidino) carbonyloxy]-1H-pyrano[3',4':6,-7]indolizino[1,2-b]quinoline-3,14-(4H,12H)-dione.

In yet another preferred embodiment of the invention HYCANTIN (Smith Kline Beecham, Philadelphia, PA) may be used in accordance with the present invention. HYCANTIN is a brand of topotecan hydrochloride for injection, a semisynthetic derivative of camptothecin. The chemical name for topotecan hydrochloride is (S)-10-[(dimethylamino)methyl]-4-

ethyl-4,8-dihydroxy-1H-pyrano[3'4':6,7] indolizino [1,2-b] quinoline-3,14-(4H, 12H)-dione monohydrochloride.

In a further embodiment of the present invention, topoisomerase inhibitors such as pentacyclic titerpenoid 5 compounds, such as  $\alpha$ -boswellic acid and its acetate,  $\beta$  boswellic acid and its acetate may be used in accordance with the present invention (U.S. Patent 5,064,823 issued Nov. 12 1991, incorporated herein by reference in its entirety).

In yet another embodiment of the present invention 10 proteins isolated from plants which are known to have topoisomerase inhibitory activity, such as GAP 31, obtainable from the seeds of *Gelonium multiflorum* and DAP 30 and DAP 32, obtainable from the leaves or seeds of *Dianthus caryophyllus*, or these proteins produced by recombinant means, are useful 15 for interfering in the interaction between human topoisomerase and HIV proteins gag and RT and therefore may be used in accordance with the present invention. The advantage of these proteins is that they demonstrate low toxicity to human cells as described in U.S. Patent No. 20 5,317,009 incorporated herein by reference in its entirety.

In addition, compounds which may also be used in accordance with the present invention, include intracellular drugs that inhibit human topo I activity - competing ligands that prevent HIV gag from interacting with human topo I - and 25 competing ligands that prevent the HIV gag and topo I complex from interacting with HIV RT.

#### 5.2.2. GENE THERAPY APPROACHES

Gene therapy approaches may also be used in accordance 30 with the present invention to inhibit the interaction of human topo I and HIV proteins. Among the compounds which may disrupt the interaction of topoisomerase with its viral targets gag and RT are antisense, ribozyme and triple helix molecules. Such molecules are designed to inhibit the 35 expression of the target gene, topoisomerase I, in HIV-infected host cells. Techniques for the production and use of antisense ribozyme and/or triple helix molecules are well



known to those of skill in the art and can be designed with respect to the cDNA (or mRNA) sequence of topo I as shown in FIG 11.

- Antisense RNA and DNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. With respect to antisense DNA, oligodeoxynucleotides derived from the translation initiation site, e.g., between the -10 and +10 regions of the target gene nucleotide sequence of interest, are preferred.
- Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. (For a review see Rossi, J., 1994, Current Biology 4:469-471). The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA, and must include the well known catalytic sequence responsible for mRNA cleavage. For this sequence see U.S. Patent No. 5,093,246, which is incorporated by reference in its entirety. As such, within the scope of the invention are engineered hammerhead motif ribozyme molecules that specifically and efficiently catalyze endonucleolytic cleavage of RNA sequences encoding target gene proteins.
- Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the molecule of interest for ribozyme cleavage sites which include the following sequences, GUA, GUU and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for predicted structural features, such as secondary structure, that may render the oligonucleotide sequence unsuitable. The suitability of candidate sequences may also be evaluated by testing their accessibility to hybridize with complementary oligonucleotides, using ribonuclease protection assays.

Nucleic acid molecules to be used in triple helix formation for the inhibition of transcription should be single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences may be pyrimidine-based, which will result in TAT and CGC triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules may be chosen that are purine-rich, for example, contain a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

Alternatively, the potential sequences that can be targeted for triple helix formation may be increased by creating a so called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

In instances wherein the antisense, ribozyme, and/or triple helix molecules described herein are utilized to inhibit mutant gene expression, it is possible that the technique may so efficiently reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by normal target gene alleles that the possibility may arise wherein the concentration of normal target gene product present may be lower than is necessary for a normal phenotype. In such cases, to ensure that substantially normal levels of target gene activity are

maintained, therefore, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal target gene activity may be introduced into cells via gene therapy methods such as those described, that do not contain  
5 sequences susceptible to whatever antisense, ribozyme, or triple helix treatments are being utilized. Alternatively, in instances whereby the target gene encodes an extracellular protein, it may be preferable to coadminister normal target gene protein in order to maintain the requisite level of  
10 target gene activity.

Anti-sense RNA and DNA, ribozyme, and triple helix molecules of the invention may be prepared by any method known in the art for the synthesis of DNA and RNA molecules. These include techniques for chemically synthesizing  
15 oligodeoxyribonucleotides and oligoribonucleotides well known in the art such as for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding the antisense RNA molecule. Such DNA  
20 sequences may be incorporated into a wide variety of vectors which incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be  
25 introduced stably into cell lines.

Various well-known modifications to the DNA molecules may be introduced as a means of increasing intracellular stability and half-life. Possible modifications include but are not limited to, the addition of flanking sequences of  
30 ribo- or deoxy-nucleotides to the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the oligodeoxyribonucleotide backbone.

35

### 5.3. THE MURINE HIV MODEL SYSTEM

The present invention relates to the expression of topo I and the co-expression of human CD4 and topo I in cell lines or in transgenic animals as a system which supports the HIV life cycle as a means of (1) identifying additional factors required to support the HIV life cycle and (2) as a system to screen test compounds, for their ability to interfere with the interaction between human topo I and HIV gag and RT. In a preferred embodiment of the invention, human topo I is expressed alone or human CD4 and topo I are co-expressed in mouse cell lines and transgenic mice.

In a further embodiment of the invention, human topo I is expressed in mouse cells or in transgenic mice. In this embodiment of the invention, internalization of the HIV virus by mouse cells is bypassed by a HIV pseudovirus. The HIV pseudovirus contains the HIV virus and an envelope protein from a virus with a natural tropism for mouse cells, such as the murine leukemia viruses (MLV). Therefore, the HIV virus is internalized by the mouse cells efficiently. These cells can then support the life cycle of the internalized HIV virus, because they express human topo I.

The present invention encompasses the expression of human topo I in animals and cells lines in order to support HIV replication. The present invention further encompasses the co-expression of human topo I and other human proteins, such as CD4, or the use of HIV pseudoviruses to promote internalization of the HIV virus.

#### 5.3.1. EXPRESSION OF HUMAN CD4 AND HUMAN TOPO I

The topo I and CD4 gene products or peptide fragments thereof, may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the topo I and CD4 gene polypeptides and peptides of the invention by expressing nucleic acid containing topo I and CD4 gene sequences are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing topo I and CD4 gene

product coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. See, for example, the techniques described in Sambrook et al., 1989, supra, and Ausubel et al., 1989, supra. Alternatively, RNA capable of encoding topo I and CD4 gene product sequences may be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis", 1984, Gait, M.J. ed., IRL Press, Oxford, which is incorporated by reference herein in its entirety.

A variety of host-expression vector systems may be utilized to express the topo I and CD4 gene coding sequences of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, exhibit the topo I and CD4 gene product of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., E. coli, B. subtilis) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing topo I and CD4 gene product coding sequences; yeast (e.g., Saccharomyces, Pichia) transformed with recombinant yeast expression vectors containing the topo I and CD4 gene product coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the topo I and CD4 gene product coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing topo I and CD4 gene product coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian

cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the topo I and CD4 gene product being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of topo I and CD4 protein or for raising antibodies to topo I and CD4 protein, for example, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which the topo I and CD4 gene product coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The topo I and CD4 gene coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of topo I and CD4 gene coding sequence will result in inactivation of the polyhedrin gene and

production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed. (E.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the topo I and CD4 gene coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing topo I and CD4 gene product in infected hosts. (E.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted topo I and CD4 gene product coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire topo I and CD4 gene, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of the topo I and CD4 gene coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription

terminators, etc. (see Bittner et al., 1987, Methods in Enzymol. 153:516-544).

5.3.2. EXPRESSION OF HUMAN TOPO I  
IN ANIMAL CELL LINES

5 The present invention encompasses the expression of human topo I and the co-expression of human CD4 and topo I in animal cell lines and their use as a system to support the HIV life cycle. In a preferred embodiment of the present invention, human CD4 and topo I are co-expressed in a mouse  
10 cell line. Host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g.  
15 cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be  
20 chosen to ensure the correct modification of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include  
25 but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3 and WI38 cell lines.

For long term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the human CD4 and/or the human  
30 topo I gene product may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription  
35 terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an



enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form 5 foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines. This method may advantageously be used to engineer cell lines which express the human CD4 or human topo I gene products. Such cell lines would be particularly 10 useful in screening and evaluation of compounds that affect the endogenous activity of the human topo I gene product.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine 15 phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk<sup>-</sup>, hgp<sup>r</sup>t<sup>-</sup> or ap<sup>r</sup>t<sup>-</sup> cells, respectively. Also, antimetabolite resistance can be used as 20 the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. 25 Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hyg<sup>r</sup>, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147).

Alternatively, any fusion protein may be readily 30 purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88: 35 8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an

amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto  $\text{Ni}^{2+}$ -nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with 5 imidazole-containing buffers.

5.3.3. EXPRESSION OF HUMAN TOPO I IN  
TRANSGENIC ANIMALS

The present invention also encompasses the expression of  
10 human topo I and the co-expression of human CD4 and topo I in transgenic animals as a model system to support the HIV life cycle. In a preferred embodiment of the invention, human CD4 and human topo I are co-expressed in the same cells in transgenic mice. In another preferred embodiment of the  
15 invention, human topo I is expressed alone in transgenic mice and a HIV pseudovirus is used to infect the animals. For example, a HIV pseudovirus which contains the HIV virus and an envelope protein from a virus with a natural tropism for murine cells, such as the murine leukemia virus (MLV), is  
20 used to bypass internalization of the HIV virus by the murine cells. These murine cells can then support the life cycle of the internalized HIV virus, because they express human topo I.

The CD4 and topo I gene products can also be expressed  
25 together in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate CD4 and topo I transgenic animals.

Any technique known in the art may be used to introduce  
30 the CD4 and topo I gene transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to pronuclear microinjection (Hoppe, P.C. and Wagner, T.E., 1989, U.S. Pat. No.  
35 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci., USA 82:6148-6152); gene targeting in embryonic stem cells

(Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 5 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

The present invention provides for transgenic animals that carry the CD4 and topo I transgene in all their cells, as well as animals which carry the transgene in some, but not 10 all their cells, i.e., mosaic animals. The transgene may be integrated as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching 15 of Lasko et al. (Lasko, M. et al., 1992, Proc. Natl. Acad. Sci. USA 89: 6232-6236). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the CD4 20 and topo I gene transgene be integrated into the chromosomal site of the endogenous CD4 or topo I gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous topo I gene are designed for the purpose of 25 integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous topo I gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous CD4 or topo I gene in only 30 that cell type, by following, for example, the teaching of Gu et al. (Gu, et al., 1994, Science 265: 103-106). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

35 In a preferred embodiment of the invention, human topo I and human CD4 are expressed in the same cells in the transgenic animals, in particular lymphocytes. For example,

human CD4 and human topo I are expressed under the same cell type specific promoter, such as the granzyme B promoter which allows expression of topo I in peripheral blood cells and in particular NK cells and T lymphocytes (Wagnier et al., 1995, 5 Mol. Cell. Biol. 92:6930-6934) or by expressing human topo I and human CD4 under a lck promoter (Allen et al. 1992, Mol. Cell. Biol. 12:2758-68; Wildin et al. 1995, J. Immunol. 155:1286-1295) which allows expression in peripheral CD4 T lymphocytes.

10 Once transgenic animals have been generated, the expression of the recombinant CD4 and topo I gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the 15 transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include but are not limited to Northern blot analysis of tissue samples obtained from the animal, in situ hybridization analysis, and RT-PCR. 20 Samples of CD4 or topo I gene-expressing tissue, may also be evaluated immunocytochemically using antibodies specific for the topo I transgene product.

25 5.4. THE USE OF MODEL SYSTEMS TO  
IDENTIFY ADDITIONAL HOST COMPONENTS  
INVOLVED IN THE HIV LIFE CYCLE

The present invention further relates to the use of cell lines and transgenic animals expressing human topo I as a screen to identify other host cellular components required to 30 support the HIV life cycle (viral entry, integration, replication and assembly). In particular, these transgenic mice have utility in screening T cell libraries to identify additional host components.

The present invention encompasses the identification of 35

additional human T cell factors required to support the HIV life cycle, using the model animal systems decried herein. These additional T cell factors which are identified may also serve as targets for intervention in order to treat HIV-5 infection. The present invention also relates to therapeutic modalities and pharmaceutical compositions designed targeting any additional T cell factors as identified herein.

A genomic library can be constructed using DNA obtained from a human T cell line. Additionally, an expression 10 library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from human T cells. The human T cells may be derived from cell culture or from a patient. In this manner, gene products made by the T cells may be expressed in cell lines and/or transgenic animals to identify 15 additional T cell components required to support the HIV life cycle.

In preparation of cDNA libraries, total RNA is isolated from human T cells. Poly(A)+ RNA is isolated from the total RNA, and cDNA prepared from the poly(A)+ RNA, all using 20 standard procedures. See, for example, Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d ed., Vol. 2 (1989). The cDNAs may be synthesized with a restriction enzyme site at their 3'-ends by using an appropriate primer and further have linkers or adaptors attached at their 5'- 25 ends to facilitate the insertion of the cDNAs into suitable cDNA cloning vectors. Alternatively, adaptors or linkers may be attached to the cDNAs after the completion of cDNA synthesis.

In preparation of genomic libraries, DNA is isolated 30 from human T cells and fragments are generated. The DNA may be cleaved at specific sites using various restriction enzymes. Alternatively, one may use DNase in the presence of manganese to fragment the DNA, or the DNA can be physically sheared, as for example, by sonication. The DNA fragments 35 can then be separated according to size by standard techniques, including but not limited to, agarose and

polyacrylamide gel electrophoresis, column chromatography and sucrose gradient centrifugation.

The genomic DNA or cDNA fragments can be inserted into suitable vectors, including but not limited to, plasmids, 5 cosmids, bacteriophages lambda or T<sub>4</sub>, and yeast artificial chromosome (YAC) [See, for example, Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989); Glover, D.M(ed.), DNA Cloning: A Practical Approach, MRL 10 Press, Ltd., Oxford, U.K., Vols. I and II (1985)].

The DNA or RNA should be at least 17 nucleotides, preferably at least 26 nucleotides, and most preferably at least 50 nucleotides in length. The nucleotide probe is hybridized under moderate stringency conditions and washed 15 under moderate, preferably high stringency conditions. Clones in libraries with insert DNA having substantial homology to the target probe will hybridize to the probe. Hybridization of the nucleotide probe to genomic or cDNA libraries is carried out using methods known in the art. One 20 of ordinary skill in the art will know that the appropriate hybridization and wash conditions depend on the length and base composition of the probe and that such conditions may be determined using standard formulae. See, for example, Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd 25 ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, Vol. 2, (1989) pp 11.45-11.57 and 15.55-15.57.

The identity of a cloned or amplified gene sequence can be verified by comparing the amino acid sequences of its three open reading frames with the amino acid sequence of 30 known gene. The identity of the cloned or amplified gene sequence may be further verified by examining its expression pattern.

Comparison of the amino acid sequences encoded by a cloned or amplified sequence may reveal that it does not 35 contain the entire gene or its promoter. In such a case the cloned or amplified gene sequence may be used as a probe to screen a genomic library for clones having inserts that

overlap the cloned or amplified gene sequence. A complete gene and its promoter may be reconstructed by splicing the overlapping gene sequences.

The ability of an additional T cell factor to enhance the HIV life cycle may be assayed in a number of ways, including but not limited to, measuring the level of HIV specific transcripts, measuring HIV RT activity, or measuring the level of HIV provirus. These assays are well known to those skilled in the art and in addition have been described *infra*. The ability of the expression of the additional T cell factor to support the HIV life cycle will be compared to those systems in which only human CD4 and topo I are expressed.

15                    5.5.    THE USE OF THE MODEL SYSTEM TO SCREEN FOR ANTI-HIV AGENTS

The present invention further relates to the model systems as described in Sections 5.3 *infra* as assay systems to screen test compounds, such as drugs, ligands (natural or synthetic), proteins, peptides and small organic molecules for their ability to interfere with the interaction between human topo I and HIV gag and RT, and therefore have utility as an anti-HIV agent.

The following assays are designed to identify compounds or compositions that interfere with the interaction between topo I and HIV proteins gag and RT. Compounds may include, but are not limited to, small molecules, peptides such as, for example, soluble peptides, including but not limited to, small molecules, peptides such as, for example, soluble peptides, including but not limited to, peptides comprising portions of topo I or the regions of topo I required for its interaction with HIV gag or RT, antibodies (including, but not limited to, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab'), and FAb expression library fragments, and epitope-binding fragments thereof), and small organic or inorganic

molecules. The compounds identified as inhibitors of the interaction between topo I and HIV proteins.

The principle of the assays to identify compounds which inhibit the interaction of topo I and HIV proteins, in particular gag and RT involves administering the compound to mouse cells stably expressing both human CD4 and topo I or to transgenic mice expressing both human CD4 and topo I infected with an appropriate concentration (i.e., TCID<sub>50</sub>) of HIV virus. Culture conditions well known to those in the art are used. A range of concentrations of the test compound may be used, in addition to a control culture wherein no test compound has been added. After incubation for an appropriate period (e.g., 7 days) of culturing, a cell-free supernatant is prepared, using standard procedures, and tested for the presence of RT activity as a measure of successful infection. The RT activity may be tested using standard techniques such as those described by, for example, Goff et al. (Goff et al., 1981, J. Virol. 38:239-248) and/or Willey et al. (Willey et al., 1988, J. Virol. 62:139-147). These references are incorporated herein by reference in their entirety.

The ability of the test compounds to inhibit HIV replication in mouse cells or transgenic mice may be assayed in a number of ways, including but not limited to, measuring the level of HIV specific transcripts, measuring HIV RT activity, measuring the level of HIV provirus or measuring the level of HIV virus. These assays are well known to those skilled in the art and in addition have been described *infra*. The ability of the test compound to inhibit the HIV life cycle will be compared to those systems in which only human CD4 and topo I are expressed.

## 5.6. PHARMACEUTICAL FORMULATIONS AND METHODS OF ADMINISTRATION

### 5.6.1. ANTIVIRALS TO BE USED IN COMBINATION WITH A TOPOISOMERASE INHIBITOR

According to the present invention, a topoisomerase I inhibitor, an inhibitor of HIV viral replication, may be used



in combination with other therapeutic agents to enhance the antiviral effect achieved. Preferably a topoisomerase I inhibitor is used in combination with another antiviral agent. Such additional antiviral agents which may be used  
5 with a topoisomerase I inhibitor include but are not limited to those which function on a different target molecule involved in viral replication, e.g., reverse transcriptase inhibitors, viral protease inhibitors, glycosylation inhibitors; those which act on a different target molecule  
10 involved in viral transmission; those which act on a different loci of the same molecule; and those which prevent or reduce the occurrence of viral resistance. One skilled in the art would know of a wide variety of antiviral therapies which exhibit the above modes of activity.

15 A topoisomerase I inhibitor can also be used in combination with retrovirus inhibitors, such as nucleoside derivatives. Nucleoside derivatives are modified forms of purine and pyrimidine nucleosides which are the building blocks of RNA and DNA. Many of the nucleoside derivatives  
20 under study as potential anti-HIV medications result in premature termination of viral DNA replication before the entire genome has been transcribed. These derivatives lack 3' substituents that can bind to subsequent nucleosides and result in chain termination. Nucleoside derivatives such as  
25 3'-azido-3'-thymidine (AZT) and dideoxyinosine (ddI) have been exploited as inhibitors of HIV-1 replication, both in vitro and in vivo. Nucleoside analogs are the currently the only licensed therapeutics for the treatment of HIV infection and AIDS (Fischl et al, 1987 N. Engl. J. Med. 317, 185-191;  
30 Mitsuya and Broder, 1987 Nature 325, 773-778). This class of compounds works by inhibiting reverse transcriptase resulting in a block in cDNA synthesis (Mitsuya and Broder, 1987), these inhibitors work early in the infectious cycle of HIV-1 and inhibit integration into T-cell genome. However, AZT  
35 therapy leads to development of resistant HIV strains (Larder 1989, 1991, *Ibid.*) and demonstrates toxicity in AIDS patients upon long-term therapy (Fischl et al., 1987, N. Engl. J.

Med. 317:185-191; Creagh-Kirk, et al., 1988, J.A.M.A. 260:3045-3048).

Further, a topoisomerase I inhibitor can be used in combination with nucleoside derivatives which include but are not limited to, 2',3'-dideoxyadenosine (ddA); 2',3'-dideoxyguanosine (ddG); 2',3'-dideoxyinosine (ddI); 2',3'-dideoxycytidine (ddC); 2',3'-dideoxythymidine (ddT); 2',3'-dideoxy-dideoxythymidine (d4T) and 3'-azido-2',3'-dideoxythymidine (AZT). Alternatively, halogenated nucleoside derivatives may be used, preferably 2',3'-dideoxy-2'-fluoronucleosides including, but not limited to, 2',3'-dideoxy-2'-fluoroadenosine; 2',3'-dideoxy-2'-fluoroinosine; 2',3'-dideoxy-2'-fluorothymidine; 2',3'-dideoxy-2'-fluorocytosine; and 2',3'-dideoxy-2',3'-didehydro-2'-fluoronucleosides including, but not limited to 2',3'-dideoxy-2',3'-didehydro-2'-fluorothymidine (Fd4T). Preferably, the 2',3'-dideoxy-2'-fluoronucleosides of the invention are those in which the fluorine linkage is in the beta configuration, including, but not limited to, 2',3'-dideoxy-2'-beta-fluoroadenosine (F-ddA), 2',3'-dideoxy-2'-beta-fluoroinosine (F-ddI), and 2',3'-dideoxy-2'-beta-fluorocytosine (F-ddC). Such combinations allow one to use a lower dose of the nucleoside derivative thus reducing the toxicity associated with that agent, without loss of antiviral activity because of the use of the topoisomerase I inhibitor. Moreover, such a combination reduces or avoids viral resistance.

Preferred combinations of topoisomerase I inhibitors and nucleoside derivatives within the scope of the present invention include an effective amount of a topoisomerase I inhibitor and an effective amount of AZT to treat HIV infection; and an effective amount of a topoisomerase I inhibitor and an effective amount of ddI.

According to the present invention, topoisomerase I inhibitors can also be used in combination with uridine phosphorylase inhibitors, including but not limited to acyclovir compounds, including benzylacyclovir (BAU);

benzyloxybenzylacyclouridine (BBAU); aminomethyl-  
benzylacyclouridine (AMBAU); aminomethyl-  
benzyloxybenzylacyclouridine (AMB-BAU); hydroxymethyl-  
benzylacyclouridine (HMBAU); and hydroxymethyl-  
5 benzyloxybenzylacyclouridine (HMBBAU).

According to the present invention, topoisomerase I  
inhibitors can be used in combination with viral protease  
inhibitors, including but not limited to, Invirase  
(saquinavir, Roche), ABT-538 (Abbott, CAS Reg. No. 155213-67-  
10 5), AG1343 (Burroughs Wellcome/Glaxo, CAS Reg. No. 161814-49-  
9). Protease inhibitors are generally thought to work  
primarily during or after assembly (*i.e.*, viral budding) to  
inhibit maturation of virions to a mature infectious state.  
For example, ABT-538 has been shown to have potent antiviral  
15 activity in vitro and favorable pharmacokinetic and safety  
profiles *in vivo* (Ho, et al., 1995, Nature 373: 123-126).  
Administration of ABT-538 to AIDS patients causes plasma HIV-  
1 levels to decrease exponentially and CD4 lymphocyte counts  
to rise substantially. The exponential decline in plasma  
20 viraemia following ABT-538 treatment reflects both the  
clearance of free virions and the loss of HIV-1 producing  
cells as the drug substantially blocks new rounds of  
infection. ABT-538 treatment reduces virus-mediated  
destruction of CD4 lymphocytes. Combining this treatment  
25 with topoisomerase I inhibitors, which inhibits at an earlier  
stage of HIV infection, viral fusion, would be likely to have  
synergistic effects and have a dramatic clinical impact.

In order to evaluate potential therapeutic efficacy of  
topoisomerase I inhibitors in combination with the antiviral  
30 therapeutics described above, these combinations may be  
tested for antiviral activity according to methods known in  
the art.

A compound of the invention can be administered to a  
human patient by itself or in pharmaceutical compositions  
35 where it is mixed with suitable carriers or excipients at  
doses to treat or ameliorate various conditions involving  
HIV-infection. A therapeutically effective dose further

refers to that amount of the compound sufficient to inhibit HIV infection. Therapeutically effective doses may be administered alone or as adjunctive therapy in combination with other treatments for HIV infection or associated diseases. Techniques for the formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, latest addition.

10

#### 5.6.2. ROUTES OF ADMINISTRATION

Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections, and optionally in a depot or sustained release formulation.

Furthermore, one may administer the agent of the present invention in a targeted drug delivery system, for example in a liposome coated with an anti-CD4 antibody. The liposomes will be targeted to and taken up selectively by cells expressing CD4.

#### 5.6.3. COMPOSITION/FORMULATION

The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, dragee-making, levitating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers, such as Hank's solution, Ringer's solution, or physiological saline buffer. For 5 transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are usually known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with 10 pharmaceutically acceptable carriers well known to those in the art.

Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral 15 ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable 20 excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium 25 carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For 30 this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be 35 added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the 5 active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or 10 liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional 15 manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, 20 e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in 25 an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or 30 continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain 35 formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of

solid hydrophobic polymers containing the therapeutic agent. Various of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, 5 release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise 10 suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

15 Many of the compounds of the invention identified as inhibitors of the interaction between topoisomerase I and the viral proteins RT and gag may be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts may be formed with many acids, including but 20 not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc.; or bases. Salts tend to be more soluble in aqueous or other protonic solvents that are the corresponding free base forms. Examples of pharmaceutically acceptable salts, carriers or excipients are 25 well known to those skilled in the art and can be found, for example, in Remington's Pharmaceutical Sciences, 18th Edition, A.R. Gennaro, Ed., Mack Publishing Co., Easton, PA, 1990. Such salts include, but are not limited to, sodium, potassium, lithium, calcium, magnesium, iron, zinc, 30 hydrochloride, hydrobromide, hydroiodide, acetate, citrate, tartrate, malate salts, and the like.



#### 5.6.4. EFFECTIVE DOSAGE

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve 5 their intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the 10 art, especially in light of the detailed disclosure provided herein.

For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. Such information can be used to 15 more accurately determine useful doses in humans.

A therapeutically effective dose refers to that amount of the compound that results in a reduction in the intensity of the infection or in amelioration of symptoms or a prolongation of survival in a patient. Toxicity and 20 therapeutic efficacy of such compounds can be determined by standard pharmaceutical, pharmacological, and toxicological procedures in cell cultures or experimental animals, e.g., for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective 25 in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD<sub>50</sub> and ED<sub>50</sub>. Compounds which exhibit high therapeutic indices are preferred. The data obtained from cell culture assays or animal studies can be 30 used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED<sub>50</sub> with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of 35 administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g.

Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p. 1).

Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are  
5 sufficient to maintain the desired modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data; e.g., the concentration necessary to achieve a 50-90% inhibition of HIV infection using the assays described herein. Dosages  
10 necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays, bioassays or immunoassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using the MEC  
15 value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%.

In cases of local administration or selective uptake,  
20 the effective local concentration of the drug may not be related to plasma concentration.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of  
25 administration and the judgment of the prescribing physician.

In immunization procedures, the amount of immunogen to be used and the immunization schedule will be determined by a physician skilled in the art and will be administered by reference to the immune response and antibody titers of the  
30 subject.

#### 5.6.5. PACKAGING

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage  
35 forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by

instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labelled for treatment of an  
5 indicated condition.

6. EXAMPLE: GAG PROTEINS ACTIVATE CELLULAR  
TOPO I IN A SPECIES SPECIFIC MANNER

The following *in vitro* experiments demonstrate that the  
10 HIV gag protein associates with human topo I in a species specific manner.

6.1. MATERIALS AND METHODS

Cell lines and plasmids.

15 HeLa cells were grown in RPMI media containing 10% fetal calf serum.

The plasmid pHXB2, which contains the entire HIV-1 provirus was used as the source of HIV viral DNA (Ratner et al., 1985, Nature 313:277-284). A 5.1-kb SacI-SalI fragment  
20 of this plasmid encoding the gag, pol, and vif genes was employed for construction of the recombinant vaccinia viruses, expression vectors, and for oligonucleotide-directed mutagenesis.

Recombinant vaccinia viruses (RVVs) expressing the  
25 entire gag polyprotein Pr55gag, capsid protein p15, and the nucleocapsid protein p24 were prepared. For Pr55gag, primer Gag-C, 5'-TTAGTTGCCCCCAAGCTTTTATTGTGACGAGGG-3' (nucleotides 1537-1669); was used to introduce a HindIII site followed by a stop codon (both underlined) using oligo-nucleotide-  
30 directed mutagenesis. After digestion with NcoI and HindIII, the fragment encoding Pr55gag was ligated to the same sites in the transfer plasmid vector pAK10. This plasmid pAKGag was used to generate RVVs using standard methods known in the art (Yasuda et al., 1990, J. Virol. 64:2788-2795; Hoshikawa  
35 et al., 1991, J. Gen. Virol. 72-2509)

Infection by RVVs and Immunoprecipitation of virus Proteins.

RVVs expressing the gag proteins and wild-type vaccinia virus were used to infect HeLa cells. At 16 hr postinfection, cells were lysed in phosphate-buffered saline containing 1% Nonidet P-40, and lysates were incubated with 5  $\mu$ l of specific antibody to either p24 or p15 for 1 hr at 4 C and subsequently with 50  $\mu$ l of Sepharose-protein G for 1 hr at 4°C. For Pr55 gag and p24, a mouse monoclonal antibody directed against p24 (NU24) was used, and for p15 a sheep polyclonal antibody directed against the carboxyl terminus of p15 was used (Weiss et al., 1992, Gene 121:203-212).

Precipitates were collected by centrifugation, washed three times with phosphate buffered saline containing 1% Nonidet P-40, and washed twice with 50 mM Tris-HCl (pH 8.0) containing 100 mM KCl/10mM MgCl<sub>2</sub>.

#### 15 Assay of topo I Activity *In Vitro*.

Relaxation assays were carried out using purified GST-Pr55gag and GST proteins and with immunoprecipitations from RVV-infected cells. GST-Pr55gag and GST were used at final concentrations of 50, 10, 2, 0.4, and 0.008  $\mu$ g/ml. Reaction mixtures containing 500 ng of supercoiled pUC19 DNA and human topo I at a final concentration of 0.1 unit per 20  $\mu$ l of Tris HCl (pH 8.0), 100 mM KCl, 10 mM MgCl<sub>2</sub>, 5 mM EDTA, and bovine serum albumin at 0.2 mg/ml. Reaction mixtures were allowed to proceed at 37°C for 15 minutes and were terminated by the addition of 5  $\mu$ l of Stop Solution [1% SDS/15% (vol/vol) glycerol/0.5% bromophenol blue]. Products were analyzed by agarose gel electrophoresis and staining with ethidium bromide.

#### 30 6.2. RESULTS

To determine if gag proteins were involved in topo I activity, HeLa cells were infected with a RVV expressing Pr55gag, and transfected with supercoiled DNA and then analyzed by Southern hybridization methods. Almost all of the transfected supercoiled DNA could be found in a relaxed form when Pr55gag was present (Fig. 1). The DNA remained predominantly in the supercoiled form when Pr55gag was

absent. To determine if the Pr55gag had topo I activity or if this could be interacting with, and perhaps enhancing, cellular topo I activity, an *in vitro* relaxation assay was employed to determine if a GST-Pr55gag fusion protein had  
5 topo I activity. It could be demonstrated that GST-Pr55gag clearly enhanced cellular topo I activity (Fig. 2). No enhancement of topo I activity was noted with GST alone.

In order to identify the HIV-1 gag proteins involved in activation of topo I, HeLa cells were infected with RVVs  
10 expressing Pr55gag, p24, and p15 proteins. Protein complexes were immunoprecipitated with specific gag antibodies, and the topo I activity present in the precipitates was analyzed. Relaxation activity was associated with only the Pr55gag and the p15 proteins, suggesting that the latter was responsible  
15 for the activation of topo I. No activity was noted in precipitates or p24 or wild-type vaccinia virus.

These results clearly show that the nucleocapsid protein p15 could activate and enhance cellular topo I activity and that the topo I activity associated with HIV-1 is of cellular  
20 origin.

#### 7. EXAMPLE: HIV RT ACTIVITY IS ENHANCED BY INTERACTION WITH CELLULAR TOPO I

The following *in vitro* experiments demonstrate that the  
25 HIV RT activity is enhanced by interacting with human topo I in a species specific manner.

##### 7.1. MATERIALS AND METHODS

###### RNA Templates.

30 Three templates were prepared. A template containing representative 5' and 3' ends of HIV RNA was prepared from plasmid pAD which was constructed as follows. PCR was used to prepare two products (AB and CD) from plasmid pHXB2 (Ratner et al., 1985, Nature 313:277-284). For AB, primers  
35 employed were A, 5'-CTGCATATAGAATTCTAATACGACTCATAGGGTCTCTC-3' (nucleotides 422-462), and B, 5'GCACTGGATCTACTCTA-3' (nucleotides 1424-1441). The former contains an EcoRI site

following by the T7 polymerase promoter (both underlined). For CD, primers employed were C, 5'CATTCGATTAGTGAACGG-3' (nucleotides 8489-8506), and D, 5'-AGATACCAGTCTAGA(TTT)<sub>8</sub>T-GAAGCACTCAAGGC-3' (nucleotides 9652-9666). Primer D contains an Xba I site (underlined) and a poly(A) region (TTT)<sub>8</sub>. Thirty cycles of denaturation for 30 sec at 94°C, primer annealing for 1 min at 52°C, and chain elongation for 1 min at 72°C were performed. Product AB was digested with EcoRI, and CD was digested with BamHI and Xba I. After digestion, AB was ligated to pUC19 that had been digested with EcoRI and Sma I. After ligation the free end of the AB product was blunt-ended with the Klenow fragment of DNA polymerase I and then ligated to the Sma I site (plasmid pAB). Product CD was directly ligated to pUC19 that had been digested with BamHI and Xba I (plasmid pCD). Plasmid pAB was digested with Pst I, blunt-ended with the Klenow fragment, and then digested with Xba I. Plasmid CD was digested with BamHI, blunt-ended with the Klenow fragment, and then digested with Xba I. The insert resulting from the latter was directly ligated to treated pAB to produce plasmid pAD. RNA templates were produced after incubation with T7 polymerase (Ambion, Austin, TX) for 90 min at 37°C with Xba I-linearized pAD. After digestion with DNase I, RNA was purified by phenol/chloroform extraction and ethanol precipitation. RNA templates containing only the 3' or 5' regions of HIV RNA were prepared from two plasmids pAD 3' and pAD 5', which were constructed as follows. To construct pAD 3' and pAD 5', which were constructed as follows. To construct pAD 3', pAD was digested with Xho I and Xba I, and the insert was directly ligated to pBluescript SKII (Stratagene). To construct pAD 5', pAD was digested with Acc I and Xba I; after Klenow treatment, the vector and insert were directly religated. RNA templates were prepared from plasmids linearized by Xba I and Xmn I digestion, respectively, and purified as above.

### Reverse Transcriptase (RT) Assays.

To analyze the influence of topo I on linear cDNA synthesis, the template containing both the 5' and 3' ends of

HIV-1 RNA was used. cDNA synthesis expected to result in a product of 1481 nucleotides was initiated using a *nef* gene primer end-labeled with [ $\theta$ - $^{32}$ P]ATP

(5'-ACCCACCTCCTCCTCCTCTTGTGC-3'; nucleotides 9002-9027).

- 5 Reaction mixtures obtained 300 ng of RNA template in 50  $\mu$ l of 50mM TrisHCL (pH 8.0) containing 150 mM KCl, 5mM MgCl $_2$ , 5mM dithiothreitol, 0.5 mM EGTA, 0.4 unit of RNasin (Promega), 0.1 unit of HIV-1 RT (Seikagaku, Kogyo, Tokyo), 100 ng of primer, 250  $\mu$ M dNTPs, bovine serum albumin at 0.1 mg/ml, GST-
- 10 p15 at 0.8  $\mu$ g/ml 10  $\mu$ l of glutathione beads (50% vol/vol; Pharmacia), and topo I at final concentrations of 0.16, 0.032 and 0.0064 unit. Samples were incubated at 37°C for 25 min. To assay the inhibitory effects of camptothecin (CPT), reactions with topo I at a final concentration of 0.16 unit
- 15 in the presence of 400, 40, 4 and 0  $\mu$ M CPT were carried out. All reactions were terminated by the addition of 25  $\mu$ l of stop solution [2% SDS/30% (vol/vol) formamide/0.1% bromophenol blue/8% Ficoll] and incubation at 85°C for 4 min. Samples were chilled on ice and electrophoresed on 1.2%
- 20 agarose gels. Gels were dried onto nylon membranes and analyzed with a Fuji BAS2000 phosphorescence imager. To analyze the effect of topo I on the synthesis and transfer of strong stop DNA (ssDNA), the two RNA templates containing the 3' and 5' ends of HIV-1 RNA were used in equal concentrations
- 25 (300 ng). cDNA synthesis was initiated using a primer corresponding to the primer binding site (PBS) (5'GTCCTGTTCGGGCGCCA-3'; nucleotides 636-653) end-labeled with [ $\theta$ - $^{32}$ P]ATP. Reaction conditions were as described above employing topo I at a final concentration of 0.16 unit/50- $\mu$ l
- 30 reaction volume. The final cDNA product would be expected to be 867 nucleotides in length. Inhibition of cDNA synthesis by CPT was carried out using final concentrations of 400, 40, and 4  $\mu$ M. Reaction products were analyzed as above.

35

## 7.2. RESULTS

These *in vitro* studies indicate that human topo I has a role in the activation of reverse transcription of HIV-1 RNA.

To investigate the role of topo I in the reverse transcription of virus RNA, the effect of topo I on both linear and elongation of newly synthesized DNA on one strand, and synthesis and strand transfer of the ssDNA. A primer  
5 corresponding to the nef region was employed to initiate cDNA synthesis, which would be expected to produce a cDNA of 1481 nucleotides (Fig. 3A). cDNA synthesis was markedly enhanced by the presence of topo I, and this effect was clearly dependent on the concentration of the enzyme. cDNA synthesis  
10 could be inhibited by CPT, a specific topo I inhibitor, also in a concentration dependent manner.

By employing a primer corresponding to the primer binding site on one RNA template to permit synthesis of the ssDNA and a second to accommodate strand transfer, cDNA  
15 synthesis was monitored. It can be seen that a cDNA product of 867 nucleotides, which consists of the 200-nucleotide ssDNA and the 667-nucleotide fragment produced after strand transfer, was synthesized in the presence of topo I (Fig. 4A). It can be seen that this could be inhibited by CPT in a  
20 concentration-dependent manner (Fig. 4B). The radioactivity evident in the lower part of the gel represents both strong stop and intermediate size cDNA products. It can be seen that the greatest levels of these are produced in reactions containing the highest concentrations of CPT, suggesting that  
25 the drug is inhibiting chain elongation rather than the strand transfer step.

#### 8. EXAMPLE: MOUSE CELLS EXPRESSING HUMAN CD4 AND TOPO I EFFECTIVELY SUPPORT HIV INFECTION

30 Human CD4 expressing mouse cells are not a natural substrate for infection with HIV. In order to show that species specificity accounts for the poor reverse transcription of the HIV genome in HIV-infected mouse cells, HIV infection was attempted in mouse cells expressing both  
35 human CD4 and topoisomerase.

##### 8.1. MATERIALS AND METHODS



The fragment of EcoRI and BamHI of human CD4 cDNA was subcloned into mammalian expression vector, pEFBOS (Mizushima et al., 1990, Nuc.Aci. Res. 18:5322). Murine cell line, L929 were transfected with human CD4 expressing vector by phosphate method. After 48 hrs, G418 (BRL) was added to 400 µg/ml. Fifteen days after selection, several colonies were examined for CD4 expression with anti-CD4 monoclonal antibody, Leu3a (Becton Dickinson). Selected colonies were again cloned and used as CD4 positive L929. This cell line were stained with Leu3a and FITC conjugated goat anti-mouse sera (Becton Dickinson) at 1:50, in the presence of 2% fetal calf serum. Relative fluorescence intensity was detected by flow cytometry.

For tagging human topo I with c-myc epitope, the sequence MEQKLISEEDL was inserted at the N-terminus of human topo I. To accomplish this the double stranded oligonucleotide 5'-GAA TTC GCC ATT ATG GAG CAG AAG CTG ATC TTC GAG GAG GAC CTG GCC ATG G GCT AGC-3' was ligated into the Sma I site of pUC19. (underlines indicate EcoRI, NcoI and Nhe I respectively) PCR was used to prepare human topo I cDNA carrying Nco I and Xba I site at its 5' and 3' site respectively, with two primers, 5'-CGT CCC TCC CCA TGG ACA TGA GTG GGG A-3' and 5'-GCC TCT TGA TCT AGA AAA CTC ATA-3' (underlines indicated Nco I and Xba I respectively). PCR products was digested with Nco I and Xba I. After digestion product was ligated to pUC-myc that had been digested with Nco I and Nhe I. After ligation, the product was digested with Eco RI and again ligated to expression vector PXMN carrying chicken actin promoter, this final expression vector was designated as pMyc-topo I. C-myc epitope tagging human topo I carries the peptide, AMD between c-myc epitope and N-terminus of human topo I. To CD4 positive L929 cells, additionally myc-human topo I expression vector, pMyc-topo I was cotransfected with hygromycin resistant vector. Transfected cells were selected with hygromycin at 300 µg/ml and G418 at 400 µg/ml. Colonies were examined with anti-myc polyclonal serum.

L929 cells were infected with HIV-1 (IIIB) by incubation for 24 hrs. and following trypsinization, cells were incubated further 7 days. Proviral DNA synthesis was analyzed by PCR employing gag(c) or U3-U5(d) specific primers. PCR using the primers for gag-amplification, SK38, 5'ATAATCCACCTATCCCAGTAGGAGAAAT3' (1544-1571) and SK39, 5'TTTGGTCCTTGTCTTATGTCCAGAATGC3' (1631-1658) or U3-U5-amplification, HL30, 5'GACAGCCTCCTAGCATTTTCGTCAC3' (265-288) and HL10, 5'AGGGTCTGAGGGATCTCTAG3' (588-607) with 0.5 unit of *Thermus aquaticus* DNA polymerase; 33 cycles of amplification were done consisting of 94°C for 30 sec., 54°C for 30 sec. and 72°C for 30 sec. beginning with 2 min. at 94°C and ending with 7 min. at 72°C. Buffer conditions included pH 8, 0, Mg<sup>2+</sup> 1.3 mM. Reaction mixture was separated by electrophoresis on 2% agarose gel and subjected to Southern hybridization. Molecular mass marker was used to determine the size of the amplified DNA. PCR products at the same region were randomly labeled with ( $\alpha$ -<sup>32</sup>P) dCTP and hybridized in Quickhybri solution (Stratagen) as recommended by manufacturer.

20

## 8.2. RESULTS

At first human CD4 was expressed in the mouse fibroblast L929 cell line. G418-selected transfectant clones were tested for their ability to bind mouse monoclonal antibody Leu3a which recognizes the gp120 binding region of the human CD4 molecule. The representative flow cytometry analyses demonstrates that expression of human CD4 on the mouse L929 cells. (Fig. 7A) Second, human topoisomerase I tagged with a c-myc epitope was expressed in the human CD4 expressing L929 cells. (Fig. 7B) Since human topo I and mouse topo I have 96% homology in amino acid sequence and 90% homology in cDNA sequence, it is impossible to distinguish them with an anti-human topo I antibody, therefore, the expression of human topo I in mouse cells was determined by Southern blot analysis with an anti-c-myc epitope sera.

To investigate the effect of human topo I on reverse transcriptase activity of the internalized HIV-1 genome, the

mouse cells were infected for 24 hours with HIV-1. Total DNA was isolated and analyzed by PCR analysis with gag primers which should only be detected after reverse transcription of the HIV-1 genome has occurred in the infected cells. At day 5 7, no provirus was detected in cells expressing human CD4 alone, however in the cells expressing both human CD4 and topo I, provirus was well amplified. Cell lysates were also analyzed by PCR with the LTR(U3-U5) primers. (Fig. 8) Therefore, these results demonstrate that the expression of 10 human topoisomerase I with human CD4 in mouse cells supports replication of HIV.

9. EXAMPLE: MOUSE CELLS EXPRESSING HUMAN  
TOPO I EFFECTIVELY SUPPORT HIV INFECTION

15 The following experiments demonstrate that once the HIV virus is internalized by mouse cells expressing human topo I, the human topo I is sufficient to support HIV replication. Internalization of the HIV virus by mouse cells expressing human topo I is bypassed by a HIV pseudovirus. 20 The HIV pseudovirus contains the HIV core virus and an envelope protein with a natural tropism for mouse cells, such as the murine leukemia viruses (MLV). The HIV-1 pseudovirus virus contains murine leukemia virus env glycoproteins and a drug resistant gene, to bypass the poor internalization of 25 RNA genome by mouse cells. After infection of pseudovirus effective reverse transcription and integration of the HIV genome is expected.

9.1. MATERIALS AND METHODS

30 Construction of HIV-hyg, HIV-bsd and pSR-E-MLV-env plasmids. The pseudovirus was produced by cotransfection of HIV-hyg or HIV-bsd with the plasmid SRE-MLV-env that expresses the envelope glycoprotein of MLV. A 1.2 kb deletion in the env gene of pNL43 the HIV-I vector was made, 35 leaving the rev responsive element and tat and rev exons intact. (Fig. 9A and Fig. 9B)

SV40 ori and hygromycine or brastcydine resistant sequences were inserted into the env deletion site. These constructs were named as HIV-hyg or HIV-BSD respectively.

The HIV-BSD was constructed as follows: a 1.2 kb deletion in the envelope gene was made from Nde I (6399) to the Bgl II site (7611). Inserted at this site was a 0.35 kb fragment including the simian virus 40 origin of replication, and a 0.8kb fragment from pBSD from Bam HI to Pst I including promoter and coding sequence for the BSD, drug resistant gene. The HIV-Hyg. was constructed as follows: the hygromycine resistant sequences were obtained from plasmid PREP4 (Invitrogen) by PCR with primers, 5'-GTC GGC CGC TCT AGA CTG CTT CAT-3' and 5'-CCT CCC CCA TCT AGA CTA TTC CTT-3'. The underlined sequence indicates Xba I site. The PCR product replaced the brastcydine resistant sequences of HIV-BSD.

The SR-E-MLV plasmid contains the MLV ecotropic env coding sequences (Donis-Keller et al., 1980, Proc. Natl. Acad. Sci. 77:1642-1645) which were subcloned into the plasmid p18MES (Fig. 9).

**METHODS.** Pseudovirus was generated by transfection of plasmid DNA into COS-1 cells by DEAE methods. 500 ng of each plasmid per 10 cm (diameter) dish of COS-1 cells was used, and 5  $\mu$ M chloroquine was added to enhance transfection efficiency. At 4 hours after transfection, the medium was replaced with Dulbecco's modified Eagle medium containing 10% fetal calf serum and antibiotics. COS-1 culture supernatants were harvested from 48 to 64 hours after transfection, filtered through a 0.22  $\mu$ m membrane, and without storage infected to L929 cells. L929 cells to be infected with pseudovirus were seeded into 6-well culture plates such that they were 60% confluent at the time of infection. Without dilution the supernatant including pseudovirus was added to the wells. Virus was allowed to absorb for 12 hrs. At 24 hrs later, cells were trypsinized and one fifth of cells per a well were spread to 10 cm dish. 2 days after hygromycine

or brastcydine was added at final concentration of 300 µg/ml, 1.5 µg/ml respectively. The medium was changed every 3 to 4 days until colonies of drug-resistant cells formed (12 to 15 days). Colonies were stained with 0.02% neutral red.

5 Pseudovirus generated by COS-1 were incubated L929 cells and selected with drugs for about 2 weeks. Since pseudovirus without env protein or drug resistant gene carrying pseudotype virus still produced colonies, the number of colonies induced pseudotype virus without env was subtracted

10 from the number of colonies induced by env-carrying pseudovirus.

## 9.2. RESULTS

The presence of colonies indicates effective

15 penetration, reverse transcription and integration of pseudovirus. The number of selected colonies were subtracted by the number of colonies post infected with pseudovirus without-env proteins as a background. See Table 1. As expected mouse cells (L929) expressing human topo I produced

20 much higher number of colonies after infection with HIV pseudovirus than mouse cells not expressing topo I.

After transfecting the HIV-hyg or HIV-bsd and the pSR-E-MLV-env plasmids in COS-1 cells, the expression of the HIV gag protein was determined by immunoblot analysis. When HIV-

25 hyg or HIV-bsd were co-transfected with pSR-E-MLV-env, bands corresponding to the processed (p24) and precursor (p55) of HIV I gag was observed. These results indicate that once the HIV virus is internalized by mouse cells expressing human topo I, the human topo I is sufficient to support HIV

30 replication.

TABLE I

5	Selected Colonies		
	Infected Cells	pNLhyg./E - pNLhyg.	pNLBS/E - pNLBS
	L929/Vector	0-20	0-10
	L929/htopo I	250	150

10. EXAMPLE: SCREENING FOR INHIBITORS OF TOPO I AS ANTI-HIV AGENTS IN MURINE CELLS

The transgenic mice and mouse cell lines expressing human topo I have utility in screening for agents that target human topo I and as a result inhibit the HIV-1 life cycle. The following example demonstrates the utility of mouse cell lines expressing human topo I in screening topo I inhibitors for their ability to inhibit HIV-1 infection. Topo I specific inhibitors, camptothecin (CPT) and TAN134A, are provided as an example of compounds to be tested for anti-HIV activity. These topo I inhibitors were analyzed for their effect on HIV reverse transcriptase activity.

10.1. MATERIALS AND METHODS

MT2 cells were incubated for 2 hours with HIV-1 IIIB strain (Ratner et al., 1985 Nature 313: 277-284) at 0.1 m.o.i. with drugs. Concentrations ranged from no drugs 0,0, AZT 0.01, 0.1, 1, 10 µg/ml, CPT 0.01, 0.1, 1 µg/ml TAN 0.01, 0.1, 1 µg/ml, CPT 0.1, 1, 10 µg/ml, TAN 0.1, 1, 10 µg/ml respectively (Fig. 5). After washing with PBS 3 times, cells were incubated again for 72 hrs with or without concentration of drugs as indicated; then lysed with 1% SDS, 10 mM Tris HCl (pH 8.0), 5 mM EDTA, 1M NaCl(Hirt's method. Following overnight incubation at 4C, large molecule of DNA were precipitated by centrifugation. Supernatants were treated with 100 µg/ml of protease K at 56° C for 2 hrs and after treated with phenol-chloroform, small molecules of DNA were precipitated and subjected to southern hybridization.

Transferred nylon membrane (Schleicher & Schuell) were probed with ( $\alpha$ - $^{32}$ P)dCTP-random-primed-labeled primer.

#### 10.2. RESULTS

5       The presence of CPT or TAN134A reduced HIV-replication in HIV-infected cells, as shown by Southern blot analysis (Fig. 5). However, TAN134A had a much more dramatic effect of reducing HIV-replication than CPT. Therefore, this example demonstrates how topoisomerase I inhibitors can be  
10 screened for their ability to inhibit the HIV life cycle.

#### 11. EXAMPLE: TRANSGENIC MICE EXPRESSING BOTH HUMAN CD4 AND HUMAN TOPO I

The following example demonstrates the successful  
15 generation of transgenic mice expressing both human CD4 and human topo I.

##### Procedure:

Fertilized eggs at early pronuclear stage, the male and female pronuclei being distinguished within the cytoplasm,  
20 were collected from the oviducts of C57 Black 6 female mice which had been mated to CBA males. Pooled zygotes were washed in fresh culture medium and stored until micromanipulation. The technique of microinjection is known to those skilled in the art and is described in U.S. Patent  
25 No. 4,873,191, incorporated herein by reference.

The plasmid expressing human topo I was prepared as follows: human topo I was expressed under a granzyme B promoter which allows expression of topo I in peripheral blood cells and in particular NK cells and T lymphocytes.  
30 The plasmid containing the granzyme B promoter (Wargnier et al., 1995, Mol. Cell. Biol. 92:6930-6934) is derived from pUC19 and is called p687. The human topo I fragment was amplified by PCR techniques, using PCR primers corresponding to the sites of the topo I fragment, the Bgl II and Bam HI.  
35 The human topo I fragment was inserted at the Bam HI/Bgl II site of p687 (Fig. 12).

The plasmid expressing human CD4 was prepared as follows: human CD4 was subcloned from plasmid pKS481/huCD4. The Bam HI to Bam HI fragment was prepared by PCR amplification. The human CD4 fragment was inserted at the 5 BamHI site of p687.

Both plasmids were linearized by digestion with Sal I prior to microinjection. The concentration of DNA used for microinjection was 250ng/ml.

After microinjection of the zygotes, there were five transfers of one cell eggs into five surrogate mothers, twenty seven, twenty, twenty, twenty and nineteen eggs. On the following day, there were three transfers of two cell embryos into three surrogate mothers, nineteen, eighteen and eighteen embryos. Eleven pups were born. Of the eleven pups six pups tested positive for expression of both the human CD4 and human topo I.

Transgenic mice expressing both human topo I and human CD4 were generated by expressing human topo I and CD4 under a lck promoter (Allen et al. 1992, Mol. Cell. Biol. 12:2758-68; Wildin et al. 1995, J. Immunol. 155:1286-1295) which allows expression in peripheral CD4 T lymphocytes. The plasmid plck-hGH contains the lck promoter (Fig. 13). The Bam HI/Bgl II fragment of the htopo I sequence (as described in Fig. 11) was inserted at the Bam HI site of plck-hGH. The hCD4 DNA was prepared by PCR amplification and inserted at the Bam HI site of plck-hGH.

12. EXAMPLE: HIV RT ACTIVITY IS  
ENHANCED IN VIVO BY CELLULAR TOPO I

The observation of species-specific enhancement of topo I activity by the gag-polyprotein raised the possibility that this activation may also be involved in the regulation of reverse transcription, and suggested that perhaps this could be important in the restriction HIV replication to cells of certain primate species. To test this hypothesis two systems were employed. In the first, attempts were made to infect mouse fibroblast cells expressing the primary receptor, CD4



and the accessory receptor fusion (Feng et al., 1996, Science 272:872-877) with and without expression of human topoisomerase I by HIV-I. In the second in attempts to overcome potential restrictions at the virus entry and  
5 initial internalization of the virus we also attempted infection of mouse fibroblasts with and without human topoisomerase expression with HIV-1 pseudovirus carrying murine leukemia virus (MLV) env glycoproteins.

10                    12.1    MATERIALS AND METHODS  
   EXPRESSION OF THE FUSION PROTEIN

   cDNAs encoding human (h) or simian (s) fusion were amplified from activated PBMC RNAs using primers hybridising to the respective 5' and 3' untranslated regions. Primers  
15 used were 5'-AGCGGT TAC CAT GGA GGG AT-3\*, and 5\*-ACA TCT GTG TTA GCT GGA GTG A-3'. Amplification products were cloned into the EcoRV site of pBIscrip (Strategene). The plasmid was then digested with Eco RI and Bam HI and the insert religated into the same site of pEFBOS (Nucl. Acid. Res.,  
20 1990, 18:5322), into which unique Eco RI and Bam HI cloning sites had been introduced. pEFBOS/hfusion or pEFBOS/sfusion was transfected into L929/hCD4 or L929/gCD4/htopo I cells together with pBabePuro. (Nucl. Acid. Res., 1990, 18:3587-3596) and then selected with puromycin at a final  
25 concentration of 3ug/ml following 2 weeks incubation. Selected colonies were infected with recombinant vaccinia virus expressing HIV-1 NL43-env gene (Aoki et al., 1991, AIDS 5:871-875) and monitored for syncytia formation under light microscopy. Cloned cells inducing syncytia were used as L929  
30 expressing the human CD4 and human fusion or simian fusion cell lines, or L929 expressing human CD4, human or simian fusion and human topoisomerase I cell lines.

   12.2    RESULTS

35                    Human CD4 and human or simian fusion molecules were expressed in the mouse fibroblast-L929 cell line. G418-selected transfectant clones were tested for their ability to

bind mouse monoclonal antibody Leu3a which recognizes the gp120 binding region of human CD4 molecule. Representative flow cytometry demonstrated high expression of human CD4 molecule. Representative flow cytometry demonstrated high expression of human CD4. In order to confirm the expression of the human or simian fusion molecules cell lines from selected colonies were infected with vaccinia virus expressing the HIV-I env gene (Aoki et al., 1991, AIDS 5:871-875) and monitored for syncytia formation. In cells expressing human fusion syncytia formation was observed. To demonstrate the expression of human topo I in the mouse cell lines antibody to a c-myc epitope was employed which had been tagged to the N terminus of the topo I molecule. This was necessary as human topo I has 96% homology in amino acid and 90% in nucleotide sequence with mouse topo I, and it is impossible to effectively distinguish them with currently available anti-human topo I antibodies. The expression of topo I was confirmed by immunoblot analysis with the anti c-myc epitope sera.

Total DNA synthesis was analyzed at day 7 using the PCR with gag primers which would be expected to detect cDNA only when effective reverse transcription in infected cells occurred (O'Brien et al., 1994, J. Virol 68:1258-1263). At day 7 in cells expressing only human CD4 and/or fusion provirus could not be detected. In contrast in cells expressing topo I, in addition to the CD4 and fusion proteins provirus synthesis was always observed. This was confirmed by additional PCR analysis using LTR specific (U3-U5) primers. Attempts to demonstrate if virus replication is complete in the CD4, fusion and topoisomerase I expressing cells were unsuccessful in that no detectable p24 antigen could be detected in cell supernatants.

The following experiments were conducted to confirm that HIV reverse transcription in vivo is enhanced by expression of human topoisomerase I. In these experiments the poor internalization of HIV-1 by murine cells was bypassed by infecting mouse cells with HIV-I pseudovirus containing MLV

ecotropic env glycoproteins. This approach permitted a determination of whether the post entry blocks to HIV-I DNA synthesis and integration can be either improved or overcome by the association of the gag proteins and human  
5 topoisomerase I. Fig. 9 shows a schematic representation of the HIV-hyg, HIV-BSD and pSR-E-MLV-env plasmids used for pseudovirus production.

Pseudovirus was produced by co-transfection of HIV-hyg or HIV-BSD with plasmid SR-E-MLV-env. Pseudovirus generated  
10 from transfection of COS-I cells were incubated with L929 cells and selected with resistant drugs following a two week incubation. L929 cells expressing human topo I showed much higher number of colonies following selection, this observation is in agreement with our findings that HIV  
15 reverse transcription is enhanced by human topoI. Similar experiments using amphotropic env expression gave the same results. Analysis of randomly selected colonies demonstrated that the cells contained HIV-I provirus of the expected size, thus showing that effective penetration, and reverse  
20 transcription and integration of pseudovirus RNA had occurred.

In order to analyze HIV-I protein synthesis, immunoblot analysis with anti p24 serum was used to demonstrate the production of gag proteins. The precursor gag polyproteins  
25 p55 and p24 could be readily detected in cells infected with the HIV-1 pseudovirus. Thus, it appears that if efficient internalization of and stabilization of the HIV-I core complex can be achieved, then the expression of human topo I effectively overcomes the restriction of HIV-I replication in  
30 mouse cells.

The results of the present study have important implications for the development of transgene animal models to study HIV-I infection. Importantly they demonstrate that expression of human topo I would appear to be necessary in  
35 the development of a successful animal model. The observation that expression of human CD4 and CXCR4 (fusion) does not result in provirus synthesis in infected murine

cells suggests that the process of internalization following attachment is extremely ineffective or that the core complex in infected mouse cells may be unstable. It is possible that additional cellular components may be required for  
5 stabilization. The identification of these and their introduction with topo I into transgenic animals should lead to a successful small animal model of HIV infection.

10 A number of references have been cited and the entire disclosures of which are incorporated herein by reference.

The present invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention,  
15 and functionally equivalent methods and components are within the scope of the invention. Indeed various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such  
20 modifications are intended to fall within the scope of the appended claims.

25

30

35

WHAT IS CLAIMED IS:

1. A method of inhibiting HIV replication in a cell infected with HIV, comprising administering an effective amount of an agent which interferes with the interaction  
5 between human topo I and HIV proteins to said cell.
2. The method of Claim 1 wherein the cell infected with HIV is in a mammal.
- 10 3. The method according to Claim 1 in which the agent is an inhibitor of human topoisomerase I activity.
4. The method according to Claim 2 in which the agent is a derivative of camptothecin.
- 15 5. A method of treating HIV infection in a subject which comprises administering an effective amount of an inhibitor of human topoisomerase I activity and an effective amount of at least one other therapeutic agent.
- 20 6. The method of Claim 5 wherein said therapeutic agent is an antiviral.
7. The method of Claim 6 wherein said antiviral is a  
25 reverse transcriptase inhibitor, a cytokine, or a cytokine inhibitor.
8. The method of Claim 6 wherein said antiviral is a nucleoside analog.
- 30 9. The method of Claim 8 wherein said nucleoside analog is AZT, ddI, ddA, d4T or 3TC.
10. The method of Claim 6 wherein said antiviral is a  
35 protease inhibitor.

11. A transgenic mouse which expresses human topoisomerase I.
12. The transgenic mouse of Claim 11 wherein the  
5 transgenic mouse further expresses human CD4 in the same cell  
as the human topoisomerase I.
13. The transgenic mouse of Claim 12 which further  
expresses fusion in the same cell as the human topoisomerase  
10 I.
14. A mouse cell line expressing human topo I.
15. The mouse cell line of Claim 14 wherein the cell  
15 line further expresses human CD4.
16. A method for identifying test compounds capable of  
inhibiting HIV infection, by (a) administering said test  
compound to the mouse cell line of Claim 14; (b) infecting  
20 the cells with a HIV pseudovirus; and (c) measuring HIV  
replication.
17. A method for identifying test compounds capable of  
inhibiting HIV infection, by (a) administering said test  
25 compound to the mouse cell line of Claim 15; (b) infecting  
the cells with HIV; and (c) measuring HIV replication.
18. A method for identifying test compounds capable of  
inhibiting HIV infection, by (a) administering said test  
30 compound to the transgenic mouse of Claim 13; (b) infecting  
the mouse with HIV; and (c) measuring HIV replication.
19. A method for identifying test compounds capable of  
inhibiting HIV infection, by (a) administering said test  
35 compound to the transgenic mouse of Claim 12; (b) infecting  
the mice with an HIV pseudovirus with a tropism for mice; and  
(c) measuring HIV replication.

1/13

COS-1  
Transfected COS-1

— 55kD  
— 24kD

FIG. 10

2/13

Homo sapiens cDNA clone E0017  
Topoisomerase I

ORIGIN

```
1  gtggcttggt gcaagaagtg ggggtgtccaa ttgagaagat ttacaacaaa acccagcgga
61  gaagtttgcc tgggccattg acatggctga tgaagactat gagttttagc cagtctcaag
121 aggcagagtt ctgtgaagag gaacagtgtg gtttgggaaa gatggataaa ctgagcctca
181 cttgccctcg tgcctcgggg agagaggcag caagtcttaa caaaccaaca tctttgcgaa
241 aagataaagc ctggagatat tataaggagg agctgagcca att
```

FIG. 11



3/13




FIG. 1

4/13




FIG. 2

5/13

A




FIG. 3A

B




FIG. 3B

6/13

A




FIG. 4A




FIG. 4B

7/13




FIG. 5

8/13

Activation of human or calf topoisomerase I  
by HIV-1 gag proteins.




FIG. 6

9/13




FIG 7A

B

L929/hCD4  
L929/hCD4/htopo I

— -100kD

FIG. 7B

10/13

A

MT2  
L929/hCD4  
L929/hCD4/htopo I

343bp -



FIG. 8A

B

MT2  
L929/hCD4  
L929/hCD4/htopo I

115bp -



FIG. 8B



11/13




FIG. 9

12/13




FIG. 12

13/13




FIG. 13

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/08686

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12Q 1/70; A01N 43/04, A61K 31/70

US CL : 435/5; 514/50, 410

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/5; 514/50, 410

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
MEDLINE, APS, WPIDS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TAKAHASHI et al. Human immunodeficiency virus type 1 reverse transcriptase: Enhancement of activity by interaction with cellular topoisomerase I. Proc. Natl. Acad. Sci. USA. June 1995, Volume 92, pages 5694-5698, see entire document.	1-3
X -- Y	US 5,422,344 A (PRIEL et al.) 06 June 1995, see entire document.	1-3 and 5-9 ----- 4 and 10
Y	FUNABASHI et al. TAN-1496 A, C and E, Diketopiperazine Antibiotics with Inhibitory Activity Against Mammalian DNA Topoisomerase I. The Journal of Antibiotics. November 1994 Volume 47, Number 11, pages 1202-1218, see entire document.	4

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

*A*	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*E*	earlier document published on or after the international filing date	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L*	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O*	document referring to an oral disclosure, use, exhibition or other means	*G*	document member of the same patent family
*P*	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search  
11 AUGUST 1997Date of mailing of the international search report  
05 SEP 1997Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Jeffrey S. Parkin, Ph.D.

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/08686

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	HO et al. Rapid turnover of plasma virions and CD4 lymphocytes in HIV-1 infection. Nature. 12 January 1995, Volume 373, pages 123-126, see entire document.	10

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/08686

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please see extra sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-10

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet(1))(July 1992)\*

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/08686

## Observations where unity of invention is lacking

This International Search Authority has found 2 inventions claimed in the International Application covered by the claims indicated below:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1

Group I, claim(s) 1-10, drawn to methods of treating or inhibiting HIV replication and infection through the administration of compounds capable of abrogating biochemical interactions between human topoisomerase I and HIV proteins.

Group II, claim(s) 11-19, drawn to transgenic murine animals and cell lines expressing human topoisomerase I, and methods employing these reagents.

and it considers that the International Application does not comply with the requirements of unity of invention (Rules 13.1, 13.2 and 13.3) for the reasons indicated below:

The inventions listed as Groups I and II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The corresponding special technical feature appears to be directed toward the identification of protein-protein interactions involving HIV-1 gag and human topoisomerase I (topo I) (refer to disclosure, page 5, lines 25-31) and the use of this interaction to identify putative inhibitors of HIV-1 viral replication. These claims are covered by the invention of Group I. However, the invention of Group II does not contain this special technical feature and is directed toward a different product (e.g., transgenic animals and cell lines expressing topo I) and method of using these products. Moreover, the invention of Group I does not employ nor require the transgenic animals and modified cell lines of Group II.

